


### Treatment Protocol

#### Step 1: Foundation




#### 1:1 dilution

Dosage:  
2 Syringes

Final Volume:  
3 ml



#### Step 2: Walls




#### 1:2 dilution

Dosage:  
2 Syringes

Final Volume:  
4.5 ml



#### Step 3: Roof



#### 1:3 dilution

Dosage:  
1-2 Syringes

Final Volume:  
6-12 ml



## Product Prep

1. Draw up appropriate amount of 1% Lidocaine or 0.9% NS into either 5 cc or 10 cc syringe
2. Attach red connector to the syringe
3. Open and attach Radiesse syringe
4. Transfer all Radiesse into syringe with Lidocaine or NS
5. Remove Radiesse syringe and attach another 5 ml or 10 ml syringe
6. Move product between two syringes at least 20 times to assure proper mixing. Be careful not to pop syringes off the connector
7. Remove one syringe and reattach Radiesse syringe
8. Remix before each use to avoid settling

## Procedure

1. Prep the entire face using Prevantics or chlorhexidine solution
2. Prior to use, clean white marker with alcohol
3. Assess and mark the patient while sitting upright
4. Avoid touching entry points with anything other than Prevantics swabs or chlorhexidine saturated gauze including while applying pressure
5. While threading and fanning, always inject retrograde with steady continuous movement
6. Continuously reassess position of the cannula tip and plane
7. Continuously reassess symmetry
8. After the procedure is finished, gently massage all the injected areas

